

PEEL BACK AT CORNER  
FOR IMPORTANT  
PRODUCT INFORMATION

Do not use if the seal is open or broken.

REORDER NO: 21013

MANUFACTURED BY:  
**Jefferson Labs**  
NORTHFIELD, MINNESOTA 55057  
1-888-215-1256  
IN 50-1419 12/2024



TAKE TIME OBSERVE LABEL  
DIRECTIONS



### Drug Facts

**Active Ingredient**  
Each 1 mL contains:  
Pyrantel base ..... Pinworm  
(as Pyrantel Pamoate) 50 mg treatment

### Indications

- For the treatment of pinworms.

### Warnings

- Abdominal cramps, nausea, vomiting, diarrhea, headache or dizziness sometimes occur after taking this drug. If any of these conditions persist, consult a physician.
- Keep this and all medications out of reach of children.
- In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.
- If you are pregnant or have liver disease, do not take this product unless directed by a physician.

### Directions

- Shake well before use.
- Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram.
- Carefully read additional product information (located on the inside of label) before using this medication. Take only according to directions and do not exceed the recommended dosage unless directed by a physician. Medication should only be taken one time as a single dose, do not repeat treatment unless directed by a physician. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See WARNINGS. If any worms other than pinworms are present before or after

### Drug Facts (continued)

#### Directions (continued)

- treatment, consult a physician. If any symptoms or pinworms are still present after treatment, consult a physician.
- This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication.

#### Dosage Information

Each mL of Pyrantel Pamoate Oral Suspension contains the equivalent of 50 mg of Pyrantel Base (as Pyrantel Pamoate).

#### Pyrantel Pamoate Oral Suspension Dosage Guide

Weight	Dosage (taken as a single dose)
Less than 25 pounds or under 2 years old	Do not use unless directed by a physician.
25 to 37 pounds	1/2 teaspoonful (125 mg)
38 to 62 pounds	1 teaspoonful (250 mg)
63 to 87 pounds	1 1/2 teaspoonfuls (375 mg)
88 to 112 pounds	2 teaspoonfuls (500 mg)
113 to 137 pounds	2 1/2 teaspoonfuls (625 mg)
138 to 162 pounds	3 teaspoonfuls (750 mg)
163 to 187 pounds	3 1/2 teaspoonfuls (875 mg)
188 pounds and over	4 teaspoonfuls (1,000 mg)

#### Other Information

**Each teaspoonful contains:** Potassium 5 mg.  
Store upright up to 25°C (77°F) with brief excursions to 40°C (104°F). DO NOT FREEZE.

#### Inactive Ingredients

Citric acid, potassium sorbate, purified water, glycerin, sodium benzoate, sucralose, vanilla custard flavor, xanthan gum.

**Questions?** 1-888-215-1256

NDC 51072-088-00

# ORAL-PRO®

## Pyrantel Pamoate Oral Suspension

### Pinworm Treatment 50 mg/mL Pyrantel Base

#### Pharmacy Bulk Pack

Keep Out of Reach of Children



## Pyrantel Pamoate Oral Suspension

READ THIS LABEL THOROUGHLY BEFORE TAKING PRODUCT

This label will tell you about pinworms and how to treat infections effectively with Pyrantel Pamoate Oral Suspension. Many who have pinworms do not have symptoms. Nevertheless, many will experience itching in and around the rectal opening. The itching may be very annoying and continued scratching can cause an irritation in this area. The itching may occur at night when sleeping. That is why restless sleep is sometimes a sign of pinworms, especially in children. It is important for you to realize that you should only treat with Pyrantel Pamoate Oral Suspension if you have seen and identified the pinworms.

Pinworms look like tiny white threads. The female pinworm is about 1/2 inch long (the male pinworm is shorter). They live in the bowel. Usually at night, the female travels to the rectal opening and lays eggs within the skin folds around the opening. This usually takes place within one to two hours after the child has been put to bed for the night. The sticky gelatin-like substance in which the eggs (they are too small to see) are deposited and the movement of the female pinworm may cause annoying itching and possibly restless sleep. By checking the rectal opening at this time, you can attempt to see the pinworm as described.

(continued) ►

The pinworm eggs normally survive for as long as three weeks. In that time, the pinworm egg may hatch and the pinworm will travel back within the rectal opening into the bowel. The pinworm reaches maturity in the bowel within two to six weeks, and the life cycle repeats itself.

Scratching will cause pinworm eggs to stick to the fingers. Reinfection will result if the fingers are placed in the mouth. The eggs, which are too small to see, contaminate whatever they come in contact with: bed clothes, underwear, toys, hands and food touched by contaminated hands. Even eggs floating in the air can be swallowed and cause infection. Eggs deposited in the environment, as stated above, can survive for as long as three weeks. Pinworms are highly contagious. Even the cleanest and most careful people can get them.

After treatment with Pyrantel Pamoate Oral Suspension, there are some rules to follow to try to prevent spreading to others and reinfection:

- Wash hands and fingernails with soap often during the day, especially before eating and after using the toilet.
- Wear tight underpants both day and night. Change them daily.
- For several days after treatment, clean the bedroom floor by vacuuming or damp mopping. Avoid dry sweeping that may stir up dust.
- After treatment, wash bed linens and night clothes (don't shake them).
- Keep the toilet seats clean.